



Monoclonal Antibodies to CD23

Technical Data Sheet

Reagent Category

Monoclonal primary antibody to CD23

Antibody Formats Supplied

□ 0.2 ml of anti CD23, Concentrate

PRODUCT# IMI14952E (Concentrate)

SOURCE: Mouse ascites fluid

IMMUNOGLOBULIN ISOTYPE: IgG

CLONE: TU1

REACTIVITY

CD23 antigen is present on a subpopulation of peripheral blood lymphocytes and in EBV-transformed lymphoblastoid cell lines. CD23 presence has been detected on neoplastic cells in B cell lymphocytic leukemias and some in centroblastic lymphoma. This antibody is suitable for use on routinely fixed paraffin embedded and frozen sections, this antibody can also be used for staining cytosmeears, cytcntrifuge preparations. This antibody is compatible with formalin fixative as well as most commonly used histological fixatives.

POSITIVE CONTROL TISSUE: Lymph node

STAINING PATTERN: Membrane

Antibody FORMAT, DILUTION and INCUBATION

This affinity-purified monoclonal antibody is provided in phosphate buffered saline pH 7.2 containing bovine serum albumin (BSA) and 0.05% sodium azide.

□ **0.2 ml of Concentrate, for paraffin section staining: Dilute 1:50 for 30-minute incubation, the dilution factors are considerably higher for 1 hour to over-night incubation.**

The dilution factor for frozen section staining and other applications are also considerably higher and must be determined by the end user for the application and the secondary staining reagents employed.

THE SUGGESTED STARTING DILUTION FACTORS ARE MERE GUIDELINES ONLY. OPTIMAL DILUTION FACTORS MUST BE DETERMINED BY THE END USER FOR THE SECONDARY STAINING REAGENTS, SPECIMEN, STAINING PROTOCOL AND APPLICATION EMPLOYED.

The above dilutions and incubation times were derived by the use of Innovex STAT-Q and HISTO-STAT staining systems. For less sensitive detection systems these dilution factors may vary and they should be determined by the end user for specific detection system employed.

CONTINUED OVER



American MasterTech

scientific laboratory supplies

POST OFFICE BOX 2539 LODI, CALIFORNIA 95241 TELEPHONE: 1 (800) 860 4073 FACSIMILE: 1 (209) 368 4136

SPECIAL USAGE GUIDANCE

For best staining results, the use of Innovex signal-Enhancing Wash Buffers (Item#: IMI00396E for Immuno-Peroxidase staining and Item#: IMI00596E for Immuno-Alkaline-phosphatase) in place of PBS or Tris buffer for the rinsing in between the incubation steps is highly recommended. The use of Innovex Enhancing Wash Buffers greatly enhances staining signals for antigenically weak or for overfixed tissues. Its use further eliminates the need for enzyme Pre-digestion, heat application (retrieval procedure) and shortens the incubation time for the primaries antibodies without addition of any extra steps to immunostaining procedures.

APPLICATIONS

This antibody is intended for use in immunolocalization of CD23 by a variety of immunoassays such as Immunocytochemistry, fluorescent labeling, ELISA, Immunoblotting and dot blotting. In immunocytochemistry, this antibody can be used in staining routinely fixed paraffin embedded or frozen histological sections for the purpose of qualitative localization of CD23. This antibody can also be used for quantitative flow cytometric assays in the indirect method.

SPECIES REACTIVITY

This antibody reactivity with human has been substantiated; other species reactivity may be characterized by the end user.

STORAGE CONDITIONS

Store in refrigerator at 2-8°C through expiration date noted on the vial.

IMPORTANT NOTE:

The interpretation of test results is the sole responsibility of the end user.

FOR IN VITRO RESEARCH USE ONLY