



## Monoclonal Antibodies to CD79a

### *Technical Data Sheet*

#### Reagent Category

Monoclonal primary antibody to CD79a

#### Antibody Formats Supplied

□ 0.2 ml of anti CD79a, Concentrate

PRODUCT# IMI17452E (Concentrate)

---

**SOURCE:** Cultured supernatant

**IMMUNOGEN:** Purified human CD79a

**IMMUNOGLOBULIN ISOTYPE:** IgG1

**CLONE:** 11E3

#### **REACTIVITY**

CD79a is one of the components of the CD79 complex which has two components CD79a and CD79b. CD79a is present from pre-B cell stage through the plasma cell stage. CD79a is present in normal and neoplastic B cells, in most acute leukemias, in B cell lymphomas and in some melomas.

This antibody is suitable for use on routinely fixed paraffin embedded and frozen sections. This antibody is compatible with formalin fixative as well as most commonly used histological fixatives.

**POSITIVE CONTROL TISSUE:** T cell Lymphoma/ Large cell lymphoma

**STAINING PATTERN:** Membrane and some Cytoplasmic staining

#### **Antibody FORMAT, DILUTION and INCUBATION**

This affinity-purified monoclonal antibody is provided in phosphate buffered saline pH 7.2 containing bovine serum albumin (BSA) and 0.05% sodium azide.

□ 0.2 ml of Concentrate, for paraffin section staining: Dilute 1:50-1:100 for 30-minute incubation, the dilution factors are considerably higher for 1 hour to over-night incubation.

**The dilution factor for frozen section staining and other applications are also considerably higher and must be determined by the end user for the application and the secondary staining reagents employed.**

THE SUGGESTED STARTING DILUTION FACTORS ARE MERE GUIDELINES ONLY. OPTIMAL DILUTION FACTORS MUST BE DETERMINED BY THE END USER FOR THE SECONDARY STAINING REAGENTS, SPECIMEN, STAINING PROTOCOL AND APPLICATION EMPLOYED.

*The above dilutions and incubation times were derived by the use of Innovex STAT-Q and HISTO-STAT staining systems. For less sensitive detection systems these dilution factors may vary and they should be determined by the end user for specific detection system employed.*

**CONTINUED OVER**



# American MasterTech

scientific laboratory supplies

POST OFFICE BOX 2539 LODI, CALIFORNIA 95241 TELEPHONE: 1 (800) 860 4073 FACSIMILE: 1 (209) 368 4136

## **SPECIAL USAGE GUIDANCE**

For best staining results, the use of Innovex signal-Enhancing Wash Buffers (Item#: IMI00396E for Immuno-Peroxidase staining and Item#: IMI00596E for Immuno-Alkaline-phosphatase) in place of PBS or Tris buffer for the rinsing in between the incubation steps is highly recommended. The use of Innovex Enhancing Wash Buffers greatly enhances staining signals for antigenically weak or for overfixed tissues. Its use further eliminates the need for enzyme Pre-digestion, heat application (retrieval procedure) and shortens the incubation time for the primaries antibodies without addition of any extra steps to immunostaining procedures.

## **APPLICATIONS**

This antibody is intended for use in immunolocalization of CD79a by a variety of immunoassays such as Immunochemistry, fluorescent labeling, ELISA, Immunoblotting and dot blotting. In immunochemistry, this antibody can be used in staining routinely fixed paraffin embedded or frozen histological sections for the purpose of qualitative localization of CD79a. This antibody can also be used for quantitative flow cytometric assays in the indirect method.

## **SPECIES REACTIVITY**

This antibody reactivity with human has been substantiated; other species reactivity may be characterized by the end user.

## **STORAGE CONDITIONS**

Store in refrigerator at 2-8°C through expiration date noted on the vial.

## **IMPORTANT NOTE:**

The interpretation of test results is the sole responsibility of the end user.

***FOR IN VITRO RESEARCH USE ONLY***