



Monoclonal Antibodies to **CD99 (Ewing sarcoma)**

Technical Data Sheet

Reagent Category

Monoclonal primary antibody to CD99
Ewing's sarcoma

Antibody Formats Supplied

□ 0.2 ml of anti CD99 -Ewing's Sarcoma Concentrate

PRODUCT# IMI17752E (Concentrate)

SOURCE: Cultured supernatant

IMMUNOGLOBULIN ISOTYPE: IgG

CLONE: 12E7

REACTIVITY

CD99, a 32kD glycoprotein is expressed on cortical thymocytes and T lymphocytes, granulose cells of the ovary, pancreatic islet cells, and sertoli cells of the testis as well as on some endothelial cells. CD99 is highly expressed on Ewing's sarcoma cells.

This antibody is suitable for use on routinely fixed paraffin embedded and frozen sections. This antibody is compatible with formalin fixative as well as most commonly used histological fixatives.

POSITIVE CONTROL TISSUE: Ewing's sarcoma

STAINING PATTERN: Membrane staining

Antibody FORMAT, DILUTION and INCUBATION

This affinity-purified monoclonal antibody is provided in phosphate buffered saline pH 7.2 containing bovine serum albumin (BSA) and 0.05% sodium azide.

□ **0.2 ml of Concentrate, for paraffin section staining: Dilute 1:10 – 1:30 for 30-minute incubation; the dilution factors are considerably higher for 1 hour to over-night incubation.**

This antibody best stains when formalin fixed paraffin embedded sections are gently retrieved with Uni-Trieve. CD antigens are heat labile and are destroyed by extreme heat. Avoid retrieving CD markers with microwave, steamer and pressure cookers.

For Background-Free and easy retrieval, Innovex Uni-Trieve (60°-70°C) universal retrieval solution is highly recommended. To retrieve with Uni-Trieve: Warm up a water bath to 60°-70°C and fill a slide holding container with Uni-Trieve solution. Place slides in a slide rack and immerse slides in a vessel or a coplin jar containing Uni-Trieve solution, bring the Uni-Trieve solution and slides to 60°-70°C and let slides incubate at 60°-70°C for 30 minutes. Rinse slides with 2 quick changes of water; No cooling period is required. Proceed with immunostaining.

The above dilutions and incubation times were derived by the use of Innovex STAT-Q and HISTO-STAT staining systems. For less sensitive detection systems these dilution factors may vary and they should be determined by the end user for specific detection system employed.

CONTINUED OVER



American MasterTech

scientific laboratory supplies

POST OFFICE BOX 2539 LODI, CALIFORNIA 95241 TELEPHONE: 1 (800) 860 4073 FACSIMILE: 1 (209) 368 4136

APPLICATIONS

This antibody is intended for use in immunolocalization of CD99 by a variety of immunoassays such as Immunocytochemistry, fluorescent labeling, ELISA, Immunoblotting and dot blotting. In immunocytochemistry, this antibody can be used in staining routinely fixed paraffin embedded or frozen histological sections for the purpose of qualitative localization of CD99. This antibody can also be used for quantitative flow cytometric assays in the indirect method.

SPECIES REACTIVITY

This antibody reactivity with human has been substantiated; other species reactivity may be characterized by the end user.

STORAGE CONDITIONS

Store in refrigerator at 2-8°C through expiration date noted on the vial.

IMPORTANT NOTE:

The interpretation of test results is the sole responsibility of the end user.

FOR PROFESSIONAL AND RESEARCH USE ONLY